

## Is local infiltration of Tranexamic Acid efficacious in reducing blood loss after simultaneous bilateral total knee Arthroplasty? A prospective randomized study

Aditya K Aggarwal

Postgraduate Institute of Medical Education and Research, India

### Extended Abstract

Key words: Arthroplasty, Blood, TKA, Arthritis, Intravenous

Abstract:

Background Total knee arthroplasty (TKA) is one of the most successful and effective surgical strategy to enhance the quality of life in patients with end-stage arthritis of knee. Significant blood loss occurring in TKA due to bone cuts, results in wound haematoma, postoperative pain, seroma formation and arthrofibrosis thus leading to impaired final outcome. TKA performed bilaterally under single anaesthesia results in substantial perioperative blood loss. Many strategies, including the use of Platelet rich plasma (PRP) and Tranexamic acid (TXA), have been employed to minimise perioperative blood loss and the resultant demand of allogenic blood transfusion [1-3]. Some researchers have reported the beneficial effects of TXA in TKA when used in an injectable form [4]. But issue of the safety of intravenous (IV) TXA has been raised by arthroplasty surgeons due to the risk of deep vein thrombosis (DVT) or pulmonary thromboembolism in high-risk patients. Even though a few studies have highlighted the efficacy of local Tranexamic Acid [2, 5, 6], it is still not very well accepted route of administration amongst the clinicians.

**Aim** of present study was to compare the efficacy of local infiltration versus intravenous administration of Tranexamic acid in reducing perioperative blood loss.

**Methods** Seventy consecutive age-, sex- and BMI-matched patients of advanced knee arthritis with tricompartmental involvement undergoing simultaneous bilateral TKA were enrolled for this prospective, randomized, double-blinded comparative study. Patients were allocated randomly by a computer generated random number table into 2 groups: Group I (IVTXA; n=35) Two doses of TXA injected intravenously (15 mg/kg 30 minutes before tourniquet deflation and repeated after 2 hours) and Group II (TTXA; n=35) Local infiltration of TXA (15 mg/kg of TXA in 100 mL of normal saline solution applied locally on to the joint surface for 10 minutes). The primary outcome measures were: total blood loss and total drain output. Blood loss was calculated from the differences between the preoperative Haemoglobin (Hb) and the lowest postoperative Hb during hospital stay or before blood transfusion. The secondary outcome measures were: number of blood units transfused and clinical and functional outcomes as evaluated by Knee Society Score (KSS), Western Ontario and McMaster Universities Arthritis Index (WOMAC) score, Visual Analogue Score (VAS) and Wound Score. Results All 70 patients were included in the study. Perioperative blood loss in group 2 ( $561.42 \pm 248.99$ ) was reduced significantly as compared to group 1 ( $1037.04 \pm 506.65$ ) with a p-value of  $<0.001$ . There was a mean reduction of blood loss by about 45% in local infiltration group as compared to intravenous group. Postoperative Hb in group 2 ( $10.30 \pm 1.11$ ) was also significantly higher as compared to group 1 ( $9.66 \pm 1.47$ ;  $p < 0.001$ ). The mean drop of haemoglobin in intravenous group was  $2.6886 \pm 1.1636$  as compared to  $1.6057 \pm 0.6791$  mg/dl in topical tranexamic acid group. This finding was statistically significant with a p-value of  $<0.001$ . Total drain output in group 2 ( $269.14 \pm 120.98$ ) was significantly reduced as compared to group 1 ( $574.14 \pm 269.03$ ;  $p < 0.001$ ). None of the patient in topical group received any allogenic blood transfusion whereas 7 patients in intravenous group received allogenic blood transfusion out of total 35 patients (20%). Thus statistically significant difference was observed in allogenic blood transfusion between the two groups ( $p = 0.000$ ). Topical tranexamic acid was more effective in maintaining higher post-operative haemoglobin, lesser need of blood transfusion and decreased total drain output as compared to intravenous tranexamic acid administration in bilateral total knee Arthroplasty with a statistically significant result.

## Research and Reviews: Orthopedics

No complication was observed in either group. Significant difference was observed in WOMAC score at 12 weeks and 6 months ( $p=0.015$ ,  $0.007$ ) and KSS score at 6 and 12 months ( $p=0.050$ ,  $0.045$ ) respectively. However no significant difference was found at 6 weeks. VAS score was significant only at 6 month with a p-value of  $0.010$  when both groups were compared. However, both the groups showed decrease in VAS score values as compared to their pre-operative values.

**Conclusion** Local infiltration of tranexamic acid significantly decreases the blood loss and improves clinical outcome following bilateral total knee arthroplasty performed under single anaesthesia and single intravenous administration.

### References

1. Iwai T, Tsuji S, Tomita T, Sugamoto K, Hideki Y, Hamada M (2013) Repeat-dose intravenous tranexamic acid further decreases blood loss in total knee arthroplasty. *Int Orthop* 37(3):441–445
2. Ishida K, Tsumura N, Kitagawa A, Hamamura S, Fukuda K, Dogaki Y, Kubo S, Matsumoto T, Matsushita T, Chin T, Iguchi T, Kurosaka M, Kuroda R (2011) Intra-articular injection of tranexamic acid reduces not only blood loss but also knee joint swelling after total knee arthroplasty. *Int Orthop* 35(11):1639–1645
3. Aggarwal AK, Shashikanth VS, Marwaha N (2014) Platelet-rich plasma prevents blood loss and pain and enhances early functional outcome after total knee arthroplasty: a prospective randomised controlled study. *Int Orthop* 38(2):387–395.
4. Good L, Peterson E, Lisander B (2003) Tranexamic acid decreases external blood loss but not hidden loss in total knee replacement. *Br J Anaesth* 90:596–599.
5. Orpen NM, Little C, Walkar G, et al (2006) Tranexamic acid reduces early postoperative blood loss after total knee arthroplasty: a prospective randomized controlled trial of 29 patients. *Knee* 13:106–110.
6. Sa-ngasoongsong P, Chankoom T, Kawinwonggowit V, et al (2011) Postoperative blood loss reduction in computer-assisted surgery total knee replacement by low dose of intra-articular tranexamic acid together with 2-hour clamp drain: a prospective triple blinded randomized controlled trial. *Orthop Rev (Pavia)* 3:e12.