

Estimation of Glycemic Index of Single and Multiple Blend Nutraceutical Biscuits

Daisy Sharma*

Department of Food Science and Nutrition, Assam down town University, Ghy, Assam

Research Article

Received: 16/07/2021

Accepted: 30/07/2021

Published: 06/08/2021

***For correspondence:**

Daisy Sharma, Department of Food Science and Nutrition, Assam down town University, Ghy, Assam

E-mail: daisysharma8991@gmail.com

Keywords: Glycemic index; Biscuits; Food; Hypoglycemic

ABSTRACT

The present study was undertaken with an aim of developing biscuits using functional ingredient like foxtail millet flour, flaxseed flour, rice bran oil and carrot flour and to assess its Glycemic Index. The GI of the biscuits was estimated through animal experimentation. The animals were divided into two groups, one group was given 0.15 mg of standard glucose and the other group was fed with equi-load of carbohydrate content present in the developed biscuit. The GI of the developed biscuit was calculated from the area under blood glucose response curve of the test food and reference food. The GI of the developed biscuit was below 47 indicating that the biscuits were low glycemic. It can therefore be concluded that the developed low glycemic index biscuits from foxtail millet, flaxseed and carrot were nutritionally rich with enhanced sensory attributes. Its hypoglycemic effect will provide potential health benefits from degenerative diseases like diabetes mellitus.

INTRODUCTION

In developing countries like India with the increasing urbanization, technological, industrial and economic advances, the demand of processed food is also increasing rapidly. Among the processed foods, bakery products, particularly biscuits command wide popularity in rural as well as urban areas among all the age groups. Biscuits are one of the most popular widely consumed processed food product in India. Foods can be fortified with nutraceuticals to enhance their nutrition as well as provide health benefits and biscuits can be excellent vehicle for fortification using nutraceuticals. Nutraceutical play a very important role in control of diabetes. According to the American Diabetes Association, (2000) if the symptoms of developing diabetes are seen, then appropriate change in diet will certainly reduce the outcome of diabetes. The increase in diabetic population of the world is among the top ten causes of death. The eating habits and patterns of the diabetic patients are the main causes for poor glycemic control.

It is reported in epidemiological studies that high fibre dietary intake can significantly reduce the incidence of diabetes. Hence the present study was planned with the following objectives:

- To develop and standardise biscuits for diabetes patients.
- To estimate the nutrient content and glycemic index of developed biscuits.

MATERIALS AND METHODS

Formulation of biscuits

The biscuits were developed from flour mix prepared by substituting wheat flour with foxtail millet flour, flaxseed flour and carrot flour at different level of incorporation. For development of single and multiple blend nutraceutical biscuits the flour mix was prepared in three different ratios viz., 80:20, 70:30, and 60:40 respectively and were used to prepare 100 gm flour mix for biscuits. For multiple blend biscuits 50% of shortening was replaced with rice bran oil. The biscuits were prepared following the guidelines outlined by (AACC) American Association of Cereal Chemists.

Acceptability trials of formulated biscuits

Acceptability of biscuits was assessed by sensory evaluation. Sensory evaluation has been defined as a scientific discipline used to evoke, measure, analyses and interpret those responses to products as perceived through the senses of sight, smell, touch, taste and hearing.

Nutrient composition of the formulated biscuits were analysed using standard methods

Glucose tolerance test

The animals used for this experiment were white albino rats obtained from College of Veterinary Sciences, Dept of Pharmacology and Toxicology, AAU Khanapara. The animals were weighed (between 180 and 220 gram) and were divided into two groups, namely group A and group B feeding on single and multiple blend biscuits(test food) respectively. Reference group was formed which was fed on standard glucose. Under group A three groups were formed namely experimental group1, experimental group 2, experimental group 3 feeding on SB1, SB2 and SB3 respectively and under group B four groups were formed namely experimental group1, experimental group 2, experimental group 3 and experimental group 4 feeding on MB1, MB2, MB3 and MB4 respectively with 5 animals in each group, notably, with no statistical differences. The rats kept in a controlled condition, 12 hours light and dark cycle. Deionized distilled water was offered ad libitum. In the use of animals for experimentation, the guidelines of the Institutional Animal Ethics Committee were followed. Formulated single and multiple blend biscuits were taken as test samples.

GTT was conducted on overnight fasted animals. Fasting blood glucose was drawn from the reference group before giving 0.15 g standard glucose dissolved in distilled water. Blood glucose was determined after feeding at 15, 30, 45, 60, 90 and 120 minutes. The same method was performed with the experimental groups by taking their fasting blood glucose and then giving them amount of test food containing 0.15 g carbohydrates for each group.

Blood samples were taken from tail tipping of the experimental animals. Blood glucose was determined by using glucose tester Device Accu check Roche.

Measurement of incremental area under curve and glycemic index

Glycemic Index (GI) for each diet was determined by calculation of Incremental Area Under two hours of blood glucose response or Curve (IAUC) for each diet and compared with the IAUC for glucose solution standard.

The glycemic index calculated for the test samples were classified by using the GI scale given below:

Glycemic index scale

Low: 55 or low

Medium: 56-69

High: 70 or more

Formulation and standardisation of biscuits

The biscuits were developed using standard recipe and acceptable level of incorporation of foxtail millet flour, flaxseed flour and carrot flour in biscuits was assessed by incorporating 20%, 30% and 40% of each flour in wheat flour and standardized for sensory characteristics. Acceptability trials were conducted by a semi-trained panel consisting of 15 numbers of judges from the Department of Food science and Nutrition and Department of Horticulture, AAU respectively. Scoring was done on nine point hedonic scale.

Moisture

The moisture contents of single and multiple blend nutraceutical biscuits are presented. The data revealed that the moisture content of single blend nutraceutical biscuits were 3.80 g/100g, 4.20 g/100g, 3.70 g/100g in SB1A, SB2B and SB3A respectively. Similarly, the moisture content of multiple blend nutraceutical biscuits were 3.41 g/100g, 3.83 g/100g, 3.15 g/100g and 4.10 g/100g in MB1A, MB2B, MB3C, and MB4A respectively. The moisture content of single and multiple blend nutraceutical biscuits increased significantly ($p < 0.05$) than control biscuits. Increase in moisture content of foods may be due to the increase in protein content because of the hydrophobic property of protein. High moisture content in biscuit may also because of higher crude fibre content which attributes to higher water absorption and moisture binding capacity.

Crude protein

The crude protein contents of multiple blend nutraceutical biscuits are presented SB1A contains highest amount of protein 9.36 g/100g, SB2B contain 8.53 g/100g and SB3A contains 5.61 g/100g. The crude protein content increased significantly ($p < 0.05$) due to incorporation of foxtail millet, flaxseed flour and decreased due to incorporation of carrot flour. Similarly the crude protein contents of multiple blend nutraceutical biscuits also significantly increased ($p < 0.05$). The data revealed that the crude protein content of multiple blend biscuits were 9.06 g/100g, 7.25 g/100g, 8.04 g/100g, and 7.74 g/100g in MB1A, MB2A, MB3A and MB4A respectively. The protein content of all single and multiple blend biscuits were found to be higher than that of control biscuits except SB3A which was prepared from incorporation of carrot flour and protein content was lower as compared with control biscuits. Addition of carrot powder lowered the total protein content of biscuit samples significantly, which might be due to lower protein content in carrot powder (0.9 g/100 g) as compared to other ingredients of composite flour blends. But a significant increase in protein content was noticed in other blends which might be due to high protein content of foxtail millet flour and flaxseed flour. Protein content of foxtail millet is 11.5 g/100g and flaxseed flour is 20 g/100g. In 2012 fortified biscuit with flaxseed and millets respectively, and reported that protein content increased because of incorporation of flaxseed and millets in biscuits and ranged between 8.62 g/100g to 10.08 g/100g.

Crude fat

The fat content of single and multiple blend nutraceutical biscuits are presented. The data revealed that the fat content of single blend nutraceutical biscuits were 14 g/100g, 18.60 g/100g and 13.23 g/100g in SB1A, SB2B and SB3A respectively. Similarly the fat contents of multiple blend nutraceutical biscuits were 15.41 g/100g, 13.06 g/100g, 14.03 g/100g, 14.33 g/100g in MB1A, MB2A, MB3A and MB4A respectively. The fat content of the developed single and multiple blend nutraceutical biscuits significantly ($p < 0.05$) decreased in all the blends than control biscuits but was at par with MB1A blend and increased significantly in SB2B. This might be due to incorporation of flaxseed in both the blend and flaxseed flour contains a high amount of fat i.e. 38 g/100g.

The biscuits made with flaxseed had a high fat content. The fat content of millet based cookies to be 13 g/100g which is similar to the findings of the present investigation. Among the different single and multiple blend nutraceutical biscuits a significant decrease ($p < 0.05$) in fat content was observed in SB3A biscuits incorporated with carrot flour when compared with control biscuits which might be due to low oil absorption capacity of carrot flour.

RESULTS AND DISCUSSION

Table 1. Nutrient composition of single and multiple blend nutraceutical biscuits.

Nutrient	Products								CD 0.05
	Control	SB1A	SB2B	SB3A	MB1A	MB2A	MB3A	MB4A	
Moisture (g)	3.56d	3.80b	4.20a	3.70c	3.41e	3.83b	3.15f	4.10a	0.11
Protein (g)	6.00g	9.36a	8.53c	5.0h	9.06b	7.25f	8.04d	7.74e	0.1
Fat (g)	14.60b	14.00c	18.60a	14.23d	15.41b	13.06d	14.03c	14.33c	0.94
Crude fibre (g)	0.82h	2.25g	16a	2.92c	3.26b	1.78e	1.70f	2.08d	0.14
Total minerals (g)	0.73f	1.78c	2.55a	1.12e	2.22b	1.40d	2.08 b	1.35d	0.14
Carbohydrate (g)	73.90a	68.7e	61.06g	73.30b	66.60f	72.21b	70.95c	70.25d	0.14
Iron (mg)	2.08e	2.22d	4.65a	2.04e	3.11b	2.31c	3.08b	2.99b	0.14
Beta-Carotene (mg)	0.024e	0.033e	0.029e	3.41a	0.033e	2.32b	2.04d	2.21c	0.17
Energy (kcal)	457.00a	437.33d	445.00b	437.00d	440.66c	437.33d	441.66c	440.66c	1.33

Carbohydrate

The carbohydrate contents of single and multiple blend nutraceutical biscuits are presented in (Table 1). The data reveal that the carbohydrate content of single blend nutraceutical was 68.70 g/100g, 61.06 g/100g and 73.30 g/100g in SB1A, SB2B and SB3A respectively. Similarly in multiple blend nutraceutical biscuits carbohydrate content was 66.6 g/100g in MB1A, 72.21 g/100g in MB2A, 70.95 g/100g in MB3A and 70.25 g/100g in MB4A. The carbohydrate contents of single and multiple blend nutraceutical biscuits decreased significantly ($p < 0.05$) than control biscuits. The high carbohydrate content in control biscuit may be due to high carbohydrate in refined wheat flour which mainly consists of endosperm portion of wheat and endosperm portion is a good source of carbohydrate.

Estimation of glycemic index of the developed single and multiple blend nutraceutical biscuits


GTT was conducted on overnight fasted animals. Fasting blood glucose was drawn from the reference group before giving 0.15 g standard glucose dissolved in distilled water. Blood glucose was determined after administration of glucose at 15, 30, 45, 60, 90 and 120 minutes. The same method was performed with the experimental groups by taking their fasting blood glucose and then giving them amount of test food containing 0.15 g carbohydrates for each group and their blood glucose level was determined in 2010.

Blood glucose response for single and multiple blend nutraceutical biscuits

The blood glucose response of each test food were evaluated and presented for single and multiple blend nutraceutical biscuits respectively. The blood glucose level was monitored at fasting period and after feeding at an interval of 15 min, 30 min, 45 min, 60 min, 90 min and 120 min, respectively. The peak blood glucose level after consumption of the biscuits was obtained in 60 min which decreased till the observation period of 120 min (2 hrs) indicating slow digestion and absorption of the nutraceutical biscuits, whereas reference food showed peak blood glucose level in 30 min and then decreased progressively till the observation period of 120 min.

This result is attributed to the presence of dietary fibre content in the different biscuits which may have protected starch from enzymatic degradation. Also, during the baking of biscuits, 1, 6 anhydro D-glucose units may have been liberated from the starch and other polysaccharides to form enzyme resistant complexes, which are not readily digested. Several authors have reported that during baking, a fraction of starch rendered itself inaccessible to amylases and total dietary fibre content increases due to retrogradation and thus slows down the digestion process.

Figure 1. Mean blood glucose response curve for different single blend nutraceutical biscuit fed to normal healthy rats.



Incremental area under blood glucose response curve (iAUC)

From the above glucose response, incremental area under blood glucose response curve was calculated. The (iAUC) value of reference food and test food are presented. The mean (iAUC) of reference food was 3202 and single blend nutraceutical biscuits were 1196, 1005 and 1508 for SB1, SB2 and SB3 respectively (Figure 1). Similarly the mean (iAUC) of reference food was 3202 and multiple blend nutraceutical biscuits were 1085, 1380, 1222, and 1335 for MB1, MB2 MB3 and MB4 respectively. From the mean (iAUC) values it can be observed that the blood glucose levels for the test products were markedly lower than orally administered glucose.

Estimation of Glycemic index of developed single and multiple blend nutraceutical biscuits

Glycemic index of single and multiple blend were calculated from their iAUC and from iAUC of reference food. The GI of test samples, single blend nutraceutical biscuits and multiple blend nutraceutical biscuits are presented.

The glycemic index of single blend nutraceutical biscuits were 37 in SB1A, 31 in SB2B and 47 in SB3A. The lowest GI was observed in SB2B blend which might be due to high content of fibre in SB2B blend due to incorporation of flaxseed flour. Similarly the glycemic index of single blend nutraceutical biscuits were 33 in MB1A, 43 in MB2A 38 in MB3A and 41 in MB4.

Findings of the present study was that the incorporation of foxtail millet in refine flour biscuits decreased the glycemic index of the biscuit which might be due to foxtail millet having higher soluble fibre and starch content. Many studies have also reported that incorporation of foxtail millet, flaxseed, buckwheat and soy flour in refine flour biscuits decreased the glycemic index of the biscuit which might be due to the high fibre content in the functional ingredients. The high soluble dietary fibre content has been found to reduce gastric emptying and absorption of glucose after a meal, resulting in improved glucose tolerance (Table 2). The soluble dietary fibre component has been reported to decrease the activity of digestive enzymes, thus resulting in incomplete hydrolysis of carbohydrates, protein and fats and delaying absorption. Soluble dietary fibres have been shown to alter food texture, structure, and viscosity, and, therefore, the rate of starch degradation and digestion, which is related to the regulation of postprandial glucose levels.

Table 2. Mean incremental area under curve (iAUC) and glycemic index for different single blend nutraceutical biscuit.

Samples	iAUC mg.min/100ml	GI
Glucose standard	3202	-
SB1A	1196	37
SB2B	1005	31
SB3A	1508	47

Table 3. Mean incremental area under curve (iAUC) and glycemic index for different multiple blend nutraceutical biscuit.

Samples	iAUC mg.min/100ml	GI
Glucose standard	3202	-
MB1A	1085	33
MB2A	1380	43
MB3A	1222	38
MB4A	1335	41

Correlation between GI and carbohydrate content of developed single and multiple blend biscuits

Pearson correlation showed that there was no significant correlation ($r=0.5859$; $p<0.01$) between the glycemic index and carbohydrate content of developed single and multiple blend biscuits (Table 3). There is little evidence about the association between GI and carbohydrate intake. Dietary glycemic index have been more strongly related to coronary heart disease and diabetes than carbohydrate intake. Total carbohydrate intake has a weak correlation with dietary glycemic index or dietary glycemic load, which takes into account the quality of carbohydrate. In 2007 also conducted a prospective cohort study, to examine the link between glycemic index (GI) and incidence of type two diabetes in an older Australian population and reported that total carbohydrate was not correlated with type two diabetes.

CONCLUSION

From the present study it can be stated that biscuits can be made with incorporation of nutraceuticals like flaxseed flour, foxtail millet flour, carrot flour without affecting the sensory attributes. These novel biscuits are nutritionally superior to that of refined wheat flour biscuits. The developed biscuits showed a low glycemic index which can be consumed by diabetic population. The outcome of the present research can be used as a valuable information for development of nutrient dense high fibre biscuits. Hence development and utilization of such functional ingredients will improve the overall health of the population as well as prevent the onset of degenerative disease and help in management of metabolic disorder like diabetes and obesity.

REFERENCES

1. Baljeet SY, Ritika BY, Reena K. Effect of incorporation of carrot pomace powder and germinated chickpea flour on the quality characteristics of biscuits. *Intern Food Res J.* 2014;21:217-222.
2. Barclay AW, Flood VM, Rochtchina E, et al. Glycemic index, dietary fiber, and risk of type 2 diabetes in a cohort of older Australians. *Diabetes Care.* 2007;30:2811-2813.
3. Brennan CS, Samyue E. Evaluation of starch degradation and textural characteristics of crude fibre enriched biscuits. *Int J Food Properties.* 2004;7:647-657.
4. Cui W, Mazza G. Physicochemical characteristics of flaxseed gum. *Food Res. Intern.* 1996;29:397-402.
5. Foster-Powell K, Holt SH, Brand-Miller JC. International table of glycemic index and glycemic load values. *Am J Clin Nutr.* 2002;76:5-56.